

ARTICLE

Open Access

The E3 ubiquitin ligase TRIM25 regulates adipocyte differentiation via proteasome-mediated degradation of PPAR γ

Jae Min Lee¹, Sun Sil Choi¹, Yo Han Lee¹, Keon Woo Khim¹, Sora Yoon¹, Byung-gyu Kim², Dougu Nam¹, Pann-Ghill Suh¹, Kyungjae Myung^{1,2} and Jang Hyun Choi¹

Abstract

Peroxisome proliferator-activated receptor gamma (PPAR γ) is a ligand-dependent transcription factor that regulates adipocyte differentiation and glucose homeostasis. The transcriptional activity of PPAR γ is regulated not only by ligands but also by post-translational modifications (PTMs). In this study, we demonstrate that a novel E3 ligase of PPAR γ , tripartite motif-containing 25 (TRIM25), directly induced the ubiquitination of PPAR γ , leading to its proteasome-dependent degradation. During adipocyte differentiation, both TRIM25 mRNA and protein expression significantly decreased and negatively correlated with the expression of PPAR γ . The stable expression of TRIM25 reduced PPAR γ protein levels and suppressed adipocyte differentiation in 3T3-L1 cells. In contrast, the specific knockdown of TRIM25 increased PPAR γ protein levels and stimulated adipocyte differentiation. Furthermore, TRIM25-knockout mouse embryonic fibroblasts (MEFs) exhibited an increased adipocyte differentiation capability compared with wild-type MEFs. Taken together, these data indicate that TRIM25 is a novel E3 ubiquitin ligase of PPAR γ and that TRIM25 is a novel target for PPAR γ -associated metabolic diseases.

Introduction

Adipose tissue plays a pivotal role in storing excess energy and is a center for energy metabolism¹. Excess body fat is considered one of the major causes of insulin resistance, dyslipidemia, type 2 diabetes, certain types of cancer, and cardiovascular disease². With respect to obesity, adipocytes exhibit an altered energy homeostasis status to not only store energy but also to generate and secrete hormones and cytokines called adipokines^{1,3}. For instance, the expression of insulin resistance-inducing adipokines, including tumor necrosis factor- α , interleukin-1, and resistin is increased in the adipose tissue of obese individuals, whereas the production of the insulin-

sensitizing hormones adiponectin or adipisin is decreased⁴. Furthermore, defects in adipocyte differentiation or function increase the probability of metabolic disorders. Thus, understanding the detailed molecular mechanisms of adipose tissue biology, especially adipocyte differentiation, may provide insights for the treatment of obesity and metabolic syndromes.

Adipocyte differentiation is tightly controlled by a series of transcription factors. A number of studies have demonstrated that peroxisome proliferator-activated receptor gamma (PPAR γ) and CCAAT/enhancer binding proteins (C/EBPs) are the primary regulators of adipogenesis and have been shown to have broad overlap in their transcriptional targets^{5,6}. Interestingly, C/EBP- α -null embryonic fibroblast cells fail to undergo adipogenesis, but this defect can be restored by the overexpression of PPAR γ ^{7,8}. Conversely, forced expression of C/EBP- α in PPAR γ -null embryonic fibroblast cells prevents the cells from differentiating⁷. These experiments demonstrated

Correspondence: Jang Hyun Choi (janghchoi@unist.ac.kr)

¹Department of Biological Sciences, Ulsan National Institute of Science and Technology (UNIST), Ulsan 689-798, Korea

²Center for Genomic Integrity (CGI), Institute for Basic Science (IBS), Department of Biological Sciences, Ulsan National Institute of Science and Technology (UNIST), Ulsan 689-798, Korea

© The Author(s) 2018



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

that PPAR γ is the key transcriptional regulator of adipogenesis and is both sufficient and necessary for fat cell differentiation.

The transcriptional activity of PPAR γ is well known to be upregulated by its ligands, such as thiazolidinediones (TZDs)⁹. PPAR γ specifically heterodimerizes with retinoid X receptor (RXR) and binds DNA repeats of the sequence AGGTCA (DR1 elements), and the PPAR γ /RXR heterodimer regulates a variety of target genes in different cells¹⁰. In the resting state (in the absence of PPAR γ ligands), PPAR γ preferentially binds to nuclear receptor corepressor 1 (NCOR1) and silencing mediator for retinoid and thyroid receptor (SMRT/NCOR2)¹¹. These complexes recruit chromatin-modifying enzymes such as histone deacetylases to repress transcription^{12,13}. However, corepressors become dissociated from PPAR γ when it is activated by ligands. Then, coactivators, such as steroid receptor coactivators, PPAR γ coactivator 1s, histone acyltransferases, and the mediator complexes, are recruited and interact with PPAR γ to promote gene transcription^{10,14,15}. In addition to ligands, post-translational modifications (PTMs), including phosphorylation, SUMOylation, acetylation, and ubiquitination, are considered as some of the major processes regulating the transcriptional activity of PPAR γ ^{16,17}. Phosphorylation of PPAR γ at Ser112 by mitogen-activated protein kinase suppresses PPAR γ transcriptional activity and adipocyte differentiation¹⁸. In contrast, while PPAR γ phosphorylation at Ser273 by cyclin-dependent kinase 5/ERK does not change its transcriptional activity, this modification has important implications for the treatment of type 2 diabetes^{19,20}. SUMOylation of PPAR γ at Lys107 blocks its transcriptional activity, while SUMOylation at Lys395 represses inflammatory gene expression by blocking nuclear factor kappa B (NF- κ B) activation^{13,21,22}. PPAR γ is also ubiquitinated and degraded in a proteasome-dependent manner^{23,24}. Together, these studies have shown that PTMs of PPAR γ are important factors in the physiological roles of PPAR γ . Thus, characterization of the novel PTMs of PPAR γ will provide important insights into our understanding of the physiological function of PPAR γ with respect to adipogenesis.

Tripartite motif (TRIM) proteins are defined as E3 ubiquitin ligases, as they contain a ring-finger domain^{25,26}. To date, more than 77 TRIM proteins have been identified in humans that are involved in a broad range of biological processes. TRIM25 (also known as EFP) is a downstream target of estrogen receptor α ²⁷. The expression of TRIM25 is upregulated in response to estrogen and is believed to mediate the effects of estrogen with respect to breast cancer as a primary response gene²⁸. In addition, TRIM25 induces the ubiquitination of retinoic acid-inducible gene 1 and regulates host antiviral innate immunity²⁹. In this study, we report a novel role of

TRIM25 in regulating metabolic pathways by mediating PPAR γ ubiquitination and its proteasome-dependent degradation, suggesting that TRIM25 may be a potential therapeutic target in PPAR γ -mediated metabolic diseases such as obesity and type 2 diabetes.

Materials and methods

Cell culture

3T3-L1, HCT116, human embryonic kidney (HEK)-293T, and HEK-293 cells were obtained from American Type Culture Collection (Manassas, VA) and cultured in Dulbecco's modified Eagle's medium (DMEM) containing fetal bovine serum (Gemini Bio-Products, West Sacramento, CA) and 1% penicillin/streptomycin (Thermo Fisher Scientific, Waltham, MA) at 37 °C under a humidified 5% CO₂ atmosphere. Adipocyte differentiation was induced by treating cells with DMEM medium containing 10% fetal bovine serum, 0.5 mM isobutylmethylxanthine, 1 μ M dexamethasone, and 850 nM insulin. Two days after the induction, the inducing medium was replaced with maintenance DMEM medium containing 10% FBS and 850 nM insulin. Lipid accumulation in the cells was detected by Oil Red O staining. Wild-type (WT) mouse embryonic fibroblast cells (MEFs) and TRIM25-knockout (KO) MEFs were kindly provided by Dr. Kyung-soo Inn (College of Pharmacy, Kyung Hee University, Korea)²⁹. All chemicals used for cell culture were obtained from MilliporeSigma (St. Louis, MO).

Plasmid constructs and RNA interference

The mammalian expression vector for FLAG-epitope tagged WT PPAR γ was described previously¹⁹. WT human TRIM25 and its deletion mutants were kindly provided by Dr. V. Narry Kim³⁰. siRNAs for TRIM25 #1 (5'-CCTCGACAAGGAAGATAAA-3') and #2 (5'-GCATCTGCTACGGAAGCAT-3') were purchased from Shanghai GenePharma (Shanghai, China). HCT116 cells were transfected with the siRNAs using Lipofectamine RNAi MAC (Thermo Fisher Scientific, Waltham, MA). The sequences used for lentiviral short hairpin RNA (shRNA) expression vectors (pLKO.1; Dharmacon, Lafayette, CO) targeting TRIM25 were #1 (5'-TTCC TCAGTTTGTACTCCAGG-3') and #2 (5'-ATGATCCA GATCTATCTTAGG-3'). For lentiviral production, HEK-293T cells were transfected with 10 μ g of the lentiviral vectors. Following infection of the cells with the viral vectors, 3T3-L1 cells were selected by incubation with 2 μ g/ml puromycin (MilliporeSigma, St. Louis, MO).

Binding assay, immunoprecipitation, and antibodies

Glutathione S-transferase (GST)-fused proteins (PPAR γ domain mutants) immobilized with glutathione-agarose were incubated with TRIM25-expressing cell lysates for 2 h at 4 °C. Protein complexes were pulled down by

centrifugation and washed four times with binding buffer. Precipitates were detected by immunoblotting using anti-GST or TRIM25 antibodies. For analyzing interactions between endogenous PPAR γ and TRIM25, 3T3-L1 adipocytes were lysed with binding buffer. Cell lysates were incubated with anti-PPAR γ or TRIM25 antibodies and analyzed by western blotting. HEK-293 cells expressing PPAR γ , TRIM25, or their mutants were lysed in binding buffer, and total cell lysates were incubated with an anti-hemagglutinin (HA) antibody at 4 °C. Immunoprecipitates or total cell lysates were analyzed with specific antibodies as indicated. The antibodies used in this study included α -TRIM25, α -PPAR γ , α -GST, α -Ub, α -aP2, and α -adipsin antibodies, which were purchased from Santa Cruz Biotechnology (Dallas, TX), while α -HA, α -actin, α -HSP90, and α -adiponectin were purchased from Cell Signaling Technology (Danvers, MA).

Identification of PPAR γ -binding complexes

PPAR γ -null MEFs⁷ were cultured in DMEM containing 10% fetal bovine serum. Nuclear extracts were prepared from PPAR γ -null MEFs that stably expressed FLAG-PPAR γ^{WT} and were incubated with the immobilized FLAG M2 agarose gel, which was then washed with binding buffer and incubated with FLAG peptide to elute the bound proteins. Immunoprecipitated proteins were separated by SDS-polyacrylamide gel electrophoresis, and specific bands were excised and digested with trypsin. The digested proteins were then analyzed by reverse-phase liquid chromatography combined with tandem mass spectrometry (LC-MS/MS) using a high-resolution hybrid mass spectrometer (LTQ-Orbitrap; Thermo Scientific, Waltham, MA). All MS/MS spectra were searched against the Uniprot protein sequence database using SequestHT (Thermo Scientific, Waltham, MA).

Reporter gene assay

HEK-293 cells were transfected with the pDR-1 luciferase reporter plasmid, PPAR γ , RXR α , and pRL-Renilla using Lipofectamine 2000 (Thermo Fisher Scientific, Waltham, MA). Following an overnight transfection, the cells were treated with rosiglitazone for 24 h. Subsequently, the cells were harvested and used to perform reporter gene assays using a Dual-Luciferase kit (Promega, Madison, WI). Luciferase activity was normalized to Renilla activity.

Gene expression analysis

Total RNA was isolated from cells or tissues using TRIzol reagent (Thermo Fisher Scientific, Waltham, MA). RNA was reverse-transcribed using an ABI Reverse Transcription Kit. Quantitative PCR reactions were performed with SYBR green fluorescent dye on an ABI9300 PCR machine (Supplementary Table 1). Relative mRNA

expression was determined using the $\Delta\Delta$ -Ct method and was normalized to tata-binding protein levels.

In vitro ubiquitination assay

FLAG-TRIM25 was transiently expressed in HEK-293 cells, purified using anti-FLAG M2 affinity gel, and eluted by adding 3 \times FLAG peptide according to the manufacturer's instructions. Recombinant GST-PPAR γ protein was incubated with 200 ng E1 (UBE1, Boston Biochem, Cambridge, MA), 500 ng E2 (UbcH5c, Boston Biochem, Cambridge, MA), 10 μ g ubiquitin, and 2 mM ATP in the absence and presence of TRIM25^{WT} or TRIM25^{CS} in 60 μ l of reaction buffer (40 mM Tris-HCl, pH 7.6, 50 mM NaCl, and 1 mM dithiothreitol) for 1 h at 37 °C. After incubating, reactions were pulled down with glutathione-agarose and washed four times with binding buffer, followed by western blotting using α -Ub or α -PPAR γ antibodies. Reaction mixtures were directly assayed by western blotting using an α -TRIM25 antibody to assess TRIM25 self-ubiquitination as a control.

Correlation analysis in human adipose tissues

TPM-normalized RNA-sequencing data from 338 subcutaneous adipose tissue samples were obtained from the GTEx portal using the sample labels downloaded from ArrayExpress³¹. Next, the Pearson correlation coefficient between TRIM25 and PPAR γ within the fat samples was calculated after removing one outlier sample whose expression in either gene was more than 5 standard deviations from the mean. The two genes showed a significant negative correlation of -0.248 (p -value $4.14\text{E-}6$).


Animals

All animal experiments were performed according to procedures approved by the Ulsan National Institute of Science and Technology's Institutional Animal Care and Use Committee. Five-week-old male C57BL/6J mice (DBL, Chungbuk, Korea) were fed a high-fat diet (60% kcal fat, D12492, Research Diets Inc., New Brunswick, NJ) for 10 weeks. The *ob/ob* mice used in this study were purchased from the Jackson Laboratory (Bar Harbor, ME).

Results

TRIM25 interacts with PPAR γ

To identify potential PTM modulators of PPAR γ , we performed proteomic analyses of binding complexes formed with PPAR γ . As shown in Fig. 1a, several bands of potential interest were observed and subjected to LC-MS/MS. Among the multiple PPAR γ -associated proteins identified, TRIM25 was of particular interest because it functions as a ubiquitin E3 ligase and as an ISG15 E3 ligase^{29,32}. TRIM25 has also been reported to mediate polyubiquitination of the DDX58 N-terminal CARD-like region, which is crucial for triggering the cytosolic signal




transduction that leads to the production of interferons in response to viral infection³³. These data led us to investigate whether TRIM25 plays a physiological role in regulating both the PTMs and the transcriptional activity of PPAR γ .

To confirm the interaction between TRIM25 and PPAR γ , PPAR γ was immunoprecipitated from cultured adipocytes and TRIM25 was detected by immunoblotting. As shown in Fig. 1b, PPAR γ was observed to interact with endogenous TRIM25. Structural aspects of the interaction between TRIM25 and PPAR γ were further investigated in vitro using recombinant PPAR γ fragments, including the A/B region (the transcriptional regulatory region), the DNA-binding domain/hinge region (DBD/H) and the ligand-binding domain. As shown in Fig. 1c, TRIM25-specifically interacted with the DBD/H domain of PPAR γ . To further identify the specific region required for TRIM25 binding to PPAR γ , smaller regions of TRIM25 were expressed into HEK-293 cells and

coimmunoprecipitated with TRIM25. As shown in Fig. 1d, amino acids 81–185 of TRIM25 were necessary for its interaction with PPAR γ .

The E3 ligase activity of TRIM25 is required to decrease PPAR γ protein stability

To elucidate the role of TRIM25 as an E3 ligase, we generated an E3 ligase-defective TRIM25 mutant by mutating cysteines 50 and 53 to serine (C50S/C53S, TRIM25^{CS}), which does not possess E3 ubiquitin ligase activity³⁴. First, we investigated the effect of TRIM25 on PPAR γ transcriptional activity. A luciferase assay was performed in HEK-293 cells expressing a peroxisome proliferator response element-containing luciferase construct. As shown in Fig. 2a, the overexpression of TRIM25^{WT} decreased PPAR γ transcriptional activity. Importantly, TRIM25^{CS} did not suppress PPAR γ transcriptional activity, indicating that the WT E3 ubiquitin ligase is important for regulating PPAR γ transcriptional




activity. Consistent with this result, specific knockdown of TRIM25 using two different small interfering RNAs (siRNA #1 and #2) increased PPAR γ transcriptional activity (Fig. 2b). Furthermore, similar effects were observed when PPAR γ was treated with rosiglitazone, a synthetic PPAR γ ligand. Two different TRIM25 siRNAs increased PPAR γ protein levels without altering mRNA levels (Fig. 2c). Together, these results suggest that TRIM25 regulates PPAR γ protein levels in a manner that is dependent on its E3 ligase activity.

After establishing that TRIM25 is an E3 ligase that regulates PPAR γ protein levels, we next investigated the effect of TRIM25 on PPAR γ protein stability. While both TRIM25^{WT} and TRIM25^{CS} interacted with PPAR γ , TRIM25^{WT} induced PPAR γ degradation, whereas TRIM25^{CS} did not (Fig. 2d). Furthermore, an increase in

TRIM25^{WT} expression resulted in decreased PPAR γ protein expression in a dose-dependent manner, which was not observed for TRIM25^{CS} (Fig. 2e), indicating that the E3 ligase activity of TRIM25 is involved in PPAR γ degradation. The overexpression of TRIM25 significantly reduced PPAR γ protein levels, which was restored by treating cells with MG132, a specific proteasome inhibitor (Fig. 2f). Together, these results strongly suggest that proteasomal degradation of PPAR γ is mediated by TRIM25.

Next, we used a cycloheximide chase assay to measure the half-life of PPAR γ in cells that overexpressed or were downregulated for TRIM25. As shown in Fig. 3a, compared with control cells, the half-life of PPAR γ was significantly reduced by the overexpression of TRIM25^{WT} but not TRIM25^{CS}. In addition, the specific knockdown of



TRIM25 could stabilize PPAR γ , suggesting that TRIM25 is a key regulator of PPAR γ protein stability (Fig. 3b).

TRIM25 is an E3 ligase that mediates the ubiquitination of PPAR γ

Next, we performed a ubiquitination analysis to further assess whether TRIM25 functions as a specific E3 ligase of PPAR γ . We used a retroviral system to generate 3T3-L1 cell lines that stably overexpressed TRIM25^{WT} or TRIM25^{CS} or carried empty vector (control cell line), and the cells were fully differentiated into adipocytes. Endogenous PPAR γ from cultured adipocytes was immunoprecipitated to assess the ubiquitination of PPAR γ by western blotting (Fig. 4a). Although the PPAR γ protein levels were decreased in cells expressing TRIM25^{WT}, TRIM25^{WT} expression resulted in increased PPAR γ ubiquitination when cells were pretreated with MG132, which was not observed following TRIM25^{CS} expression. Consistent with these results, transiently expressed TRIM25^{WT} enhanced PPAR γ ubiquitination compared with the TRIM25^{CS} mutant in HEK-293 cells (Fig. 4b). To further confirm the direct effect of TRIM25 on the ubiquitination of PPAR γ , we performed an *in vitro* ubiquitination assay using purified TRIM25 and PPAR γ . As expected, TRIM25 induced the ubiquitination of recombinant PPAR γ , while TRIM25^{CS} did not (Fig. 4c). These results suggest that PPAR γ is a substrate of TRIM25.


TRIM25 is negatively correlated with PPAR γ

In the present study, our results demonstrated that TRIM25 regulates PPAR γ protein stability. Thus, we next investigated the physiological role of TRIM25 in adipocyte differentiation. First, we examined the physiological relevance of TRIM25 and PPAR γ during adipocyte differentiation. Western blot and real-time PCR analyses

revealed that TRIM25 was expressed in pre-adipocytes and that its expression was significantly decreased during adipogenesis (Fig. 5a, b). In contrast with TRIM25 expression, the expression of PPAR γ and adipogenic marker genes, including aP2 and adiponectin, were significantly increased, indicating that TRIM25 expression is negatively correlated with that of PPAR γ . Next, we measured TRIM25 expression in adipose tissue from high-fat-diet (HFD)-induced obesity model mice and genetically induced obesity model mice (*ob/ob*). As shown in Fig. 5c, d, TRIM25 expression was significantly downregulated in adipose tissue from both HFD and *ob/ob* mice compared with control mice, whereas PPAR γ expression was dramatically upregulated. To examine whether the correlation between TRIM25 and PPAR γ expression in adipose tissue is relevant in humans, we used RNA-sequencing data of 338 human adipose tissue samples from the Genotype-Tissue Expression (GTEx) project to analyze the correlation between TRIM25 and PPAR γ gene expression^{35,36}. As shown in Fig. 5e, the expression of TRIM25 was significantly negative correlated with that of PPAR γ . Taken together, these results suggest that the expression of TRIM25 is inversely correlated with that of PPAR γ in both mice and humans.

TRIM25 suppresses adipocyte differentiation

PPAR γ is both necessary and sufficient for adipogenesis, and alterations in PPAR γ activity affect adipogenesis. Consistent with the observed TRIM25-mediated downregulation of PPAR γ stability, when TRIM25^{WT} was expressed in differentiating 3T3-L1 cells, triglyceride accumulation was suppressed, as confirmed by Oil Red O staining (Fig. 6a). Furthermore, the protein and mRNA expression levels of adipocyte-selective genes, including aP2, C/EBP- α , adiponectin, adipisin, glucose transporter-4



(GLUT4), and LPL were also reduced (Fig. 6b, c). However, the TRIM25^{CS} mutant did not alter the expression of adipogenic markers. In addition, the specific knockdown of TRIM25 using a lentivirus expressing a shRNA targeting TRIM25 enhanced adipocyte differentiation (Fig. 6d) and the expression of adipocyte-specific proteins and genes compared with that of the control (Fig. 6e, f).


To further confirm the effect of TRIM25 on adipogenesis, we used WT and TRIM25 KO MEFs. Compared with WT MEFs, KO MEFs exhibited dramatically enhanced lipid accumulation (Fig. 7a). Furthermore, the expression of adipogenic markers was significantly increased in KO MEFs compared with WT MEFs (Fig. 7b, c). Taken together, these results suggest that TRIM25 plays a key role in the regulation of PPAR γ -dependent adipogenic processes.

Discussion

Adipose tissue is the center of systemic metabolic regulation in rodents and humans. The dysregulation of adipocyte differentiation and/or physiological function

results in metabolic disorders such as type 2 diabetes^{37,38}. As PPAR γ functions as a master regulator of fat cell differentiation and glucose/lipid metabolism, understanding the mechanisms of PPAR γ regulation is crucial for combating metabolic disorders. In this study, we demonstrated that the E3 ubiquitin ligase TRIM25 regulates the protein stability of PPAR γ , inhibiting adipogenesis in both 3T3-L1 cells and MEFs. Several lines of evidence support this conclusion. First, TRIM25 directly interacts with PPAR γ and increases ubiquitin- and proteasome-mediated PPAR γ degradation. Second, specific knockdown of TRIM25 dramatically increases adipocyte differentiation. Finally, the expression of TRIM25 is inversely correlated with that of PPAR γ in adipose tissue in both mice and humans.

Although numerous studies have reported on the regulation of PPAR γ expression at the transcriptional level, the regulation of PPAR γ expression at the protein level has been poorly studied. Recently, multiple lines of evidence have suggested that PPAR γ is regulated by PTMs, including phosphorylation, SUMOylation, and



ubiquitination^{17,24}. In particular, the polyubiquitination of PPAR γ is crucial for regulating both its expression and transcriptional activity. MKRN1, an E3 ligase of PPAR γ , was observed to induce PPAR γ ubiquitination and its proteasome-mediated degradation³⁹. However, the ubiquitination of PPAR γ does not always increase its degradation. In a report by Kilroy et al.⁴⁰, TZDs increased the expression of E3 ligase, Siah2, and Siah2, which are required for adipogenesis by ubiquitinating PPAR γ . In addition, NEDD4, an E3 ubiquitin ligase, interacts with and ubiquitinates PPAR γ , which results in increased PPAR γ protein stability by inhibiting its proteasomal degradation⁴¹. In this study, we demonstrated that the ubiquitination of PPAR γ by a novel E3 ligase, TRIM25, promoted PPAR γ proteasomal degradation and suppressed adipogenesis. These results suggest that there are several distinct pathways for ubiquitin-mediated PPAR γ regulation and that each E3 ligase regulates PPAR γ

stability in response to different cellular signals or conditions in adipocytes.

Recent reports have suggested that the protein stability of PPAR γ can be regulated by phosphorylation. Keshet et al.⁴² demonstrated that c-Abl-mediated tyrosine phosphorylation of PPAR γ increases its accumulation. In addition, epidermal growth factor receptor has been suggested to induce PPAR γ phosphorylation, which could lead to MDM2 E3 ubiquitin ligase-mediated PPAR γ ubiquitination and degradation^{43,44}. Thus, we tested whether known phosphorylation sites of PPAR γ could modulate TRIM25-mediated PPAR γ ubiquitination and degradation. TRIM25 reduced the stability of the phospho-defective mutants PPAR γ S112A and PPAR γ S273A (data not shown)^{18,19}. Interestingly, phosphorylation at Y78 of PPAR γ dramatically suppressed TRIM25-mediated PPAR γ ubiquitination and increased PPAR γ accumulation (Supplementary





Figure 1). Tyrosine phosphorylation of PPAR γ at Y78 is induced by the c-Abl or c-Src kinases, both of which are required for fat cell differentiation^{42,45,46}. Therefore, it is likely that the modulation of PPAR γ phosphorylation at Y78 is an important regulatory mechanism for its protein stability and fat cell differentiation activity. Further mechanistic studies on the role of Y78 phosphorylation in TRIM25-mediated PPAR γ regulation should be conducted in the future.

In this study, we demonstrated that TRIM25 binds to PPAR γ in adipocytes and that the expression of TRIM25 and PPAR γ is inversely correlated in adipocytes in both mice and humans (Fig. 5). Furthermore, we demonstrated the expression of TRIM25 during adipogenic processes, and both TRIM25 gene and the protein levels were transiently reduced after inducing adipocyte differentiation (Fig. 5). The exact molecular mechanism of how the expression of TRIM25 is reduced during adipogenic differentiation process remains unclear. However, we believe that the decrease in TRIM25-mediated degradation of PPAR γ during differentiation, in combination with the increase of PPAR γ gene expression and increased activity

of PPAR γ , directly induces the expression of PPAR γ itself or C/EBP α during differentiation. In addition to adipocytes, TRIM25 and PPAR γ are also expressed in different cells and tissues, especially tumors^{47–49}. Multiple lines of evidence suggest that PPAR γ acts as a tumor suppressor because it plays a role in inflammation and glucose metabolism in cancer. The overexpression of PPAR γ suppresses cell survival by inhibiting cell proliferation and tumor growth⁵⁰. In contrast, the silencing of PPAR γ in cancer has the opposite effect on the activation of the cancer survival pathway⁵¹. TRIM25 has also been linked to several cancers. In particular, TRIM25 is overexpressed in breast, colorectal, and lung cancers, while PPAR γ is downregulated in these cancers compared with normal tissues^{52–54}. These results suggest that TRIM25 may act as a crucial regulator of PPAR γ in different types of tumors. Although further studies are needed to elucidate the molecular relationship between TRIM25 and PPAR γ in tumor progression, our results extend our knowledge regarding the regulation of PPAR γ and potential targets for the development of novel therapeutic approaches targeting obesity and metabolic diseases.



Acknowledgements

This work was supported by the Korea Mouse Phenotyping Project (2016M3A9D5A01952411), a National Research Foundation of Korea (NRF) grant (2017R1A2B2007378 and 2014M3A9D8034464) funded by the Korean government (MSIP) and an NRF grant (2016R1A6A3A11932338) funded by the Ministry of Education.

Author details

¹Department of Biological Sciences, Ulsan National Institute of Science and Technology (UNIST), Ulsan 689-798, Korea. ²Center for Genomic Integrity (CGI), Institute for Basic Science (IBS), Department of Biological Sciences, Ulsan National Institute of Science and Technology (UNIST), Ulsan 689-798, Korea

Conflict of interest

The authors declare that they have no conflict of interest.

Publisher's note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Supplementary information accompanies this paper at <https://doi.org/10.1038/s12276-018-0162-6>.

Received: 14 March 2018 Revised: 29 June 2018 Accepted: 4 July 2018
Published online: 15 October 2018

References

- Rosen, E. D. & Spiegelman, B. M. Adipocytes as regulators of energy balance and glucose homeostasis. *Nature* **444**, 847–853 (2006).
- Must, A. et al. The disease burden associated with overweight and obesity. *JAMA* **282**, 1523–1529 (1999).
- Jung, U. J. & Choi, M. S. Obesity and its metabolic complications: the role of adipokines and the relationship between obesity, inflammation, insulin resistance, dyslipidemia and nonalcoholic fatty liver disease. *Int. J. Mol. Sci.* **15**, 6184–6223 (2014).
- Ouchi, N., Parker, J. L., Lugus, J. J. & Walsh, K. Adipokines in inflammation and metabolic disease. *Nat. Rev. Immunol.* **11**, 85–97 (2011).
- Lemberger, T., Desvergne, B. & Wahli, W. Peroxisome proliferator-activated receptors: a nuclear receptor signaling pathway in lipid physiology. *Ann. Rev. Cell Dev. Biol.* **12**, 335–363 (1996).
- Rosen, E. D. & Spiegelman, B. M. Molecular regulation of adipogenesis. *Ann. Rev. Cell Dev. Biol.* **16**, 145–171 (2000).
- Rosen, E. D. et al. C/EBP α induces adipogenesis through PPAR γ : a unified pathway. *Genes Dev.* **16**, 22–26 (2002).
- Wu, Z. et al. Cross-regulation of C/EBP α and PPAR γ controls the transcriptional pathway of adipogenesis and insulin sensitivity. *Mol. Cell* **3**, 151–158 (1999).
- Lehmann, J. M. et al. An antidiabetic thiazolidinedione is a high affinity ligand for peroxisome proliferator-activated receptor γ (PPAR γ). *J. Biol. Chem.* **270**, 12953–12956 (1995).
- DiRenzo, J. et al. Peroxisome proliferator-activated receptors and retinoic acid receptors differentially control the interactions of retinoid X receptor heterodimers with ligands, coactivators, and corepressors. *Mol. Cell. Biol.* **17**, 2166–2176 (1997).

11. Wagner, B. L., Norris, J. D., Knotts, T. A., Weigel, N. L. & McDonnell, D. P. The nuclear corepressors NCoR and SMRT are key regulators of both ligand- and 8-bromo-cyclic AMP-dependent transcriptional activity of the human progesterone receptor. *Mol. Cell. Biol.* **18**, 1369–1378 (1998).
12. Fajas, L. et al. The retinoblastoma-histone deacetylase 3 complex inhibits PPAR γ and adipocyte differentiation. *Dev. Cell* **3**, 903–910 (2002).
13. Picard, F. et al. Sirt1 promotes fat mobilization in white adipocytes by repressing PPAR- γ . *Nature* **429**, 771–776 (2004).
14. Puigserver, P. et al. A cold-inducible coactivator of nuclear receptors linked to adaptive thermogenesis. *Cell* **92**, 829–839 (1998).
15. Yuan, C.-X., Ito, M., Fondell, J. D., Fu, Z.-Y. & Roeder, R. G. The TRAP220 component of a thyroid hormone receptor-associated protein (TRAP) coactivator complex interacts directly with nuclear receptors in a ligand-dependent fashion. *Proc. Natl Acad. Sci. USA* **95**, 7939–7944 (1998).
16. Katsura, S., Okumura, T., Ito, R., Sugawara, A. & Yokoyama, A. Identification of posttranslational modifications in peroxisome proliferator-activated receptor γ using mass spectrometry. *PPAR Res.* **2014**, 468925 (2014).
17. Choi, S.-S., Park, J. & Choi, J. H. Revisiting PPAR γ as a target for the treatment of metabolic disorders. *BMB Rep.* **47**, 599 (2014).
18. Hu, E., Kim, J. B., Sarraf, P. & Spiegelman, B. M. Inhibition of adipogenesis through MAP kinase-mediated phosphorylation of PPAR gamma. *Science* **274**, 2100 (1996).
19. Choi, J. H. et al. Anti-diabetic drugs inhibit obesity-linked phosphorylation of PPAR [ggr] by Cdk5. *Nature* **466**, 451–456 (2010).
20. Choi, J. H. et al. Antidiabetic actions of a non-agonist PPAR [ggr] ligand blocking Cdk5-mediated phosphorylation. *Nature* **477**, 477–481 (2011).
21. Floyd, Z. E. & Stephens, J. M. Control of peroxisome proliferator-activated receptor γ 2 stability and activity by SUMOylation. *Obesity* **12**, 921–928 (2004).
22. Ohshima, T., Koga, H. & Shimotohno, K. Transcriptional activity of peroxisome proliferator-activated receptor γ is modulated by SUMO-1 modification. *J. Biol. Chem.* **279**, 29551–29557 (2004).
23. Hauser, S. et al. Degradation of the peroxisome proliferator-activated receptor γ is linked to ligand-dependent activation. *J. Biol. Chem.* **275**, 18527–18533 (2000).
24. Beekun, O., Fleskens, V. & Kalkhoven, E. Posttranslational modifications of PPAR- γ : fine-tuning the metabolic master regulator. *Obesity* **17**, 213–219 (2009).
25. Raymond, A. et al. The tripartite motif family identifies cell compartments. *EMBO J.* **20**, 2140–2151 (2001).
26. Watanabe, M. & Hatakeyama, S. TRIM proteins and diseases. *J. Biochem.* **161**, 135–144 (2017).
27. Orimo, A., Inoue, S., Ikeda, K., Noji, S. & Muramatsu, M. Molecular cloning, structure, and expression of mouse estrogen-responsive finger protein Efp. Co-localization with estrogen receptor mRNA in target organs. *J. Biol. Chem.* **270**, 24406–24413 (1995).
28. Ikeda, K., Orimo, A., Higashi, Y., Muramatsu, M. & Inoue, S. Efp as a primary estrogen-responsive gene in human breast cancer. *FEBS Lett.* **472**, 9–13 (2000).
29. Gack, M. U. et al. TRIM25 RING-finger E3 ubiquitin ligase is essential for RIG-I-mediated antiviral activity. *Nature* **446**, 916 (2007).
30. Kwon, S. C. et al. The RNA-binding protein repertoire of embryonic stem cells. *Nat. Struct. Mol. Biol.* **20**, 1122 (2013).
31. Rocca-Serra, P. et al. ArrayExpress: a public database of gene expression data at EBI. *C. R. Biol.* **326**, 1075–1078 (2003).
32. Zou, W. & Zhang, D.-E. The interferon-inducible ubiquitin-protein isopeptide ligase (E3) EFP also functions as an ISG15 E3 ligase. *J. Biol. Chem.* **281**, 3989–3994 (2006).
33. Oshiumi, H., Matsumoto, M., Hatakeyama, S. & Seya, T. Riplet/RNF135, a RING finger protein, ubiquitinates RIG-I to promote interferon- β induction during the early phase of viral infection. *J. Biol. Chem.* **284**, 807–817 (2009).
34. Meroni, G. & Diez-Roux, G. TRIM/RBCC, a novel class of 'single protein RING finger'E3 ubiquitin ligases. *Bioessays* **27**, 1147–1157 (2005).
35. Consortium, G. The Genotype-Tissue Expression (GTEx) pilot analysis: multi-tissue gene regulation in humans. *Science* **348**, 648–660 (2015).
36. Lonsdale, J. et al. The genotype-tissue expression (GTEx) project. *Nat. Genet.* **45**, 580 (2013).
37. Hotamisligil, G. S. Inflammation and metabolic disorders. *Nature* **444**, 860 (2006).
38. Trujillo, M. E. & Scherer, P. E. Adipose tissue-derived factors: impact on health and disease. *Endocr. Rev.* **27**, 762–778 (2006).
39. Kim, J. et al. Suppression of PPAR γ through MKRN1-mediated ubiquitination and degradation prevents adipocyte differentiation. *Cell Death Differ.* **21**, 594 (2014).
40. Kilroy, G., Kirk-Ballard, H., Carter, L. E. & Floyd, Z. E. The ubiquitin ligase Siah2 regulates PPAR γ activity in adipocytes. *Endocrinology* **153**, 1206–1218 (2012).
41. Li, J. J. et al. Ubiquitin ligase NEDD4 regulates PPAR γ stability and adipocyte differentiation in 3T3-L1 cells. *Sci. Rep.* **6**, 38550 (2016).
42. Keshet, R. et al. c-Abl tyrosine kinase promotes adipocyte differentiation by targeting PPAR-gamma 2. *Proc. Natl Acad. Sci. USA* **111**, 16365–16370 (2014).
43. Xu, Y. et al. EGFR/MDM2 signaling promotes NF- κ B activation via PPAR γ degradation. *Carcinogenesis* **37**, 215–222 (2016).
44. Shi, J. et al. Pioglitazone inhibits EGFR/MDM2 signaling-mediated PPAR γ degradation. *Eur. J. Pharmacol.* **791**, 316–321 (2016).
45. Choi, S. et al. Novel phosphorylation of PPAR γ ameliorates obesity-induced adipose tissue inflammation and improves insulin sensitivity. *Cell. Signal.* **27**, 2488–2495 (2015).
46. Sun, Y. et al. Requirement of SRC-family tyrosine kinases in fat accumulation. *Biochemistry* **44**, 14455–14462 (2005).
47. Han, S. & Roman, J. Peroxisome proliferator-activated receptor γ : a novel target for cancer therapeutics? *Anticancer Drugs* **18**, 237–244 (2007).
48. Tontonoz, P. & Spiegelman, B. M. Fat and beyond: the diverse biology of PPAR γ . *Annu. Rev. Biochem.* **77**, 289–312 (2008).
49. Heikel, G., Choudhury, N. R. & Michlewski, G. The role of Trim25 in development, disease and RNA metabolism. *Biochem. Soc. Trans.* **44**, 1045–1050 (2016).
50. Tontonoz, P. et al. Terminal differentiation of human liposarcoma cells induced by ligands for peroxisome proliferator-activated receptor γ and the retinoid X receptor. *Proc. Natl Acad. Sci. USA* **94**, 237–241 (1997).
51. Gimun, G. D. et al. APC-dependent suppression of colon carcinogenesis by PPAR γ . *Proc. Natl Acad. Sci. USA* **99**, 13771–13776 (2002).
52. Walsh, L. A. et al. An integrated systems biology approach identifies TRIM25 as a key determinant of breast cancer metastasis. *Cell Rep.* **20**, 1623–1640 (2017).
53. Sun, N., Xue, Y., Dai, T., Li, X. & Zheng, N. Tripartite motif containing 25 promotes proliferation and invasion of colorectal cancer cells through TGF- β signaling. *Biosci. Rep.* **37**, BSR20170805 (2017).
54. Qin, X., Qiu, F. & Zou, Z. TRIM25 is associated with cisplatin resistance in non-small-cell lung carcinoma A549 cell line via downregulation of 14-3-3 α . *Biochem. Biophys. Res. Commun.* **493**, 568–572 (2017).