

Home Search Collections Journals About Contact us My IOPscience

Strong terahertz emission from electromagnetic diffusion near cutoff in plasma

This content has been downloaded from IOPscience. Please scroll down to see the full text. 2015 New J. Phys. 17 043045 (http://iopscience.iop.org/1367-2630/17/4/043045)

View the table of contents for this issue, or go to the journal homepage for more

Download details:

IP Address: 114.70.7.203 This content was downloaded on 19/10/2015 at 03:33

Please note that terms and conditions apply.

New Journal of Physics

The open access journal at the forefront of physics

Deutsche Physikalische Gesellschaft **OPG**

IOP Institute of Physics

Published in partnership with: Deutsche Physikalische Gesellschaft and the Institute of Physics

PAPER

CrossMark

OPEN ACCESS

RECEIVED 7 November 2014

REVISED 3 March 2015

ACCEPTED FOR PUBLICATION 19 March 2015

PUBLISHED 22 April 2015

Content from this work may be used under the terms of the Creative Commons Attribution 3.0 licence.

Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.



Strong terahertz emission from electromagnetic diffusion near cutoff in plasma

M-H Cho^{1,5}, Y-K Kim², H Suk^{3,7}, B Ersfeld⁴, D A Jaroszynski^{4,7} and M S Hur^{1,6}

¹ School of Natural Science, UNIST, Ulsan, 689-798, Korea

- ² School of Electrical and Computer Engineering, UNIST, Ulsan, 689-798, Korea
- ^b Department of Physics and Photon Science, GIST, Gwangju, 500-712, Korea
- Department of Physics, Scottish Universities Physics Alliance and University of Strathclyde, Glasgow G4 0NG, UK
- Center for Relativistic Laser Science, Institute for Basic Science (IBS), Gwangju 500-712, Korea
- Author to whom any correspondence should be addressed.

⁷ Co-corresponding authors.

E-mail: mshur@unist.ac.kr, D.A.Jaroszynski@strath.ac.uk and hysuk@gist.ac.kr

Keywords: laser-plasma, terahertz radiation, laser wakefield

Abstract

A new mechanism for electromagnetic emission in the terahertz (THz) frequency regime from laserplasma interactions is described. A localized and long-lasting transverse current is produced by two counter-propagating short laser pulses in weakly magnetized plasma. We show that the electromagnetic wave radiating from this current source, even though its frequency is close to cut-off of the ambient plasma, grows and diffuses towards the plasma-vacuum boundary, emitting a strong *monochromatic* THz wave. With driving laser pulses of moderate power, the THz wave has a field strength of tens of MV m⁻¹, a frequency of a few THz and a *quasi*-continuous power that exceeds all previous *monochromatic* THz sources. The novelty of the mechanism lies in a diffusing electromagnetic wave close to cut-off, which is modelled by a continuously driven complex diffusion equation.

1. Introduction

Far-infrared light has been expected to be uniquely advantageous for probing and imaging the structure and dynamics of matter compared with other light sources. However, the characteristics of the radiation sources in that frequency regime have not been satisfactory with respect to various requirements. Enormous efforts are now being made to fill the perceived terahertz (THz) gap with the development of high power THz sources because of their potential high-impact applications in science and technology. Diverse sources have been developed using both electronic and optical methods, for example the laser-plasma-based schemes [1–20], which promise compact THz sources with notably high field amplitude over a wide range of frequencies, and conventional beam driven vacuum sources such as gyrotrons, backward wave oscillators etc.

Linear mode conversion and two-colour schemes are known to yield short-duration and wide-band THz pulses with remarkably high amplitude. The latter is especially attractive because a high THz amplitude up to 2 GV m^{-1} can be expected [21]. In this scheme, a slow component of a strong photocurrent is efficiently yielded by phase-controlled photoionization at each laser field extremum [22, 23], which is essential in generating a broad-band THz pulse.

While the peak power of these wide-band THz sources is increasing rapidly, the average power of available *narrow-band* THz sources remains very low. A *monochromatic* THz source based on laser-plasma interactions and wakefields is feasible [6]. By converting the longitudinal current of the wakefield to a transverse one using an external magnetic field, this Cherenkov wake scheme extends the pulse duration and increases the emission power [18–20]. However, the field amplitude or power from such a monochromatic THz source is still low compared with wide-band schemes. Considering the numerous important applications such as high-contrast



imaging [24–26] or interferometry of Tokamak plasmas [27], it is necessary to provide suitable *monochromatic* sources that have high tunability, compactness, and above all, high power.

In this paper, we propose a new mechanism of generating high-intensity monochromatic THz radiation from laser-plasma interactions. The fundamental idea is to evoke a strong, localized, long-lasting electron oscillation in plasma, which acts as a radiating antenna emitting a continuous THz wave. In practice, the radiation current source can be generated by the strong ponderomotive force from two short laser pulses colliding at a desired position inside the plasma [28]. The current generated in this way is maintained for a long time by the plasma oscillation mechanism even after the passage of the driving laser pulses. The longitudinal current induced in this way oscillates electrostatically. To obtain electromagnetic radiation, i.e. the THz wave, some fraction of the longitudinal electron oscillation is converted to a transverse one by a weak transverse external magnetic field. Because the radiation from this current source oscillates at the plasma frequency, the electromagnetic fields are at the cut-off of the ambient plasma and cannot propagate under normal conditions, which is indeed the very reason why plasma oscillations cannot be easily converted to transverse electromagnetic waves in laser-plasma systems. However, we have discovered that if the electromagnetic wave is constantly driven near cut-off, as in our case, the field grows as its energy is fed by the driving current. Furthermore, both electric and magnetic components of the field diffuse strongly into the plasma. As a consequence, the diffusing electromagnetic field eventually propagates across the plasma-vacuum boundary and is converted to an electromagnetic wave emitted into free space. This concept of THz generation is shown schematically in figure 1. Our scheme shares a common feature with the Cherenkov wake driven by a single pulse in using the external magnetic field to convert a longitudinal oscillation to a transverse one. However, the stronger, local electron oscillation by colliding pulses, and the diffusion-growth mechanism are two unique features of our scheme.

As the plasma frequency can be easily controlled in the few-THz regime, this mechanism can be utilized as a new THz radiation source. As we will show later through numerical simulations, the resultant amplitude of the THz emission reaches tens of MV m⁻¹ and has a sub-nanosecond pulse duration. Such high amplitudes in the regime of narrow-band THz pulses with long pulse duration are not readily obtained with conventional methods in compact systems. Note that from wide-band THz sources, such as the two-colour scheme, short THz pulses with peak amplitudes higher by two orders of magnitude are available [21].

This paper is organised as follows. In section 2, we describe the diffusion and growth of the electromagnetic field near cutoff theoretically and using simulations. In section 3, a theoretical scaling law for the THz emission amplitude as a function of the amplitude of the driving pulse is derived. Then, in section 4 we give a summary and conclusions.

2. Electromagnetic diffusion and growth near cutoff

The diffusion of the electromagnetic field, which is a key factor for enhanced THz emission in our scheme, is apparent in the field evolution modelled by a constantly driven complex diffusion equation. Such diffusion of the electromagnetic wave is quite different from the well-known magnetic field diffusion into conducting material, where just the slowly varying magnetic component diffuses while the electric component remains negligibly small. To describe electromagnetic diffusion near cut-off, we start from the wave equation:

$$\frac{\partial^2 E_y}{\partial x^2} - \frac{1}{c^2} \frac{\partial^2 E_y}{\partial t^2} = \frac{\omega_p^2}{c^2} E_y + \mu_0 \frac{\partial J_y}{\partial t}.$$
(1)

Similar equations describing the field evolution in the two-colour scheme can be found in [29]. Equation (1) differs from those in so far as, firstly, J_v originates from the ponderomotive force of the laser pulses rather than





from photoionization, and secondly, the role of the laser pulses in our scheme is limited to an initial source current J_y . Thus E_y in equation (1) includes just the THz field without the laser field. Note that the first term on the right-hand side is the self-current induced by E_y itself, while the second term is a constant driving current. Assuming that the electric field amplitude evolves slowly, the electric field E_y can be written as $\hat{E}e^{-i\omega t}$, where \hat{E} is a slowly evolving function of time. Since $\omega \simeq \omega_p$, i.e. E_y is nearly at cut-off, the first term on the right-hand side cancels out the second time derivative of $e^{-i\omega t}$ on the left-hand side. Neglecting the second time derivative of \hat{E} in accordance with the slowly varying assumption, we obtain a constantly driven diffusion-like equation of the electric field:

$$\frac{\partial^2 \hat{E}}{\partial x^2} + \frac{2i\omega_p}{c^2} \frac{\partial \hat{E}}{\partial t} = -i\omega_p \mu_0 J_0, \tag{2}$$

where J_0 is the current amplitude defined by $J_y = J_0 e^{-i\omega_p t}$. The transverse current J_y is induced by the external magnetic field (B_0) from the longitudinal current J_x , which is driven by the beat of two counter-propagating laser pulses. Because the pulse duration is very short, a few tens of femtoseconds, the longitudinal current J_x is generated almost instantaneously at the moment of pulse collision. Once J_x is formed, it continues to oscillate by the regular plasma oscillation mechanism. Since the conversion rate from J_x to J_y is very small for $\omega_c = eB_0/m \ll \omega_p$, J_x acts as feeding J_y constantly. Thus J_0 can be considered temporally quasi-constant, leading to $\partial_t J_y \simeq -i\omega_p J_0 e^{-i\omega_p t}$. The THz pulse generated by such a constantly oscillating current is expected to have a long pulse duration, as will be confirmed later. Note that equation (2) takes the familiar form of a driven diffusion equation, except that the diffusion coefficient, which is $-ic^2/2\omega_p$, is imaginary.

By applying Laplace and inverse Laplace transforms to equation (2), the solution can be represented in integral form as follows:

$$\frac{e\hat{E}(x,t)}{mc\omega_p} = \frac{i+1}{8\pi} \int_{-\infty}^{\infty} dx' j_0 e^{-y^2/\bar{\sigma}_J^2} \int_{\gamma-i\infty}^{\gamma+i\infty} ds \frac{e^{i\eta\gamma+s\bar{s}}}{s^{3/2}},$$
(3)

where the variables are normalized as $y = \omega_p |x - x'|/c$, $\bar{\sigma}_J = \omega_p \sigma_J/c$, $\eta = \sqrt{2is}$, $\bar{t} = \omega_p t$, and $j_0 = J_0 / en_0 c$. In deriving equation (3) we have assumed that the transverse current amplitude J_0 takes on a very narrow Gaussian form, $J_0 = \hat{J}_0 e^{-x^2/\sigma_J^2}$. Here σ_J is the spatial length of the current source, which corresponds to half the laser pulse duration.

Approximate solutions of equation (3) can be obtained in certain limits. First, for $y/\bar{t} \gg 1$, i.e. far from the current source and temporally in the early stage, applying the steepest descent method leads to

$$\frac{e\hat{E}\left(x,\,t\right)}{mc\omega_{p}} \simeq -\frac{\bar{\sigma}_{J}j_{0}}{\sqrt{2}}\frac{\bar{t}^{3/2}}{\bar{x}^{2}}\cos\left(\frac{\bar{x}^{2}}{2\bar{t}}+\frac{\pi}{4}\right),\tag{4}$$

with $\bar{x} = \omega_p x/c$. Second, for x = 0, i.e. at the oscillation centre of the current, by applying Watson's lemma to the first order term we obtain an asymptotic form as a function of *t* as follows:

$$\frac{e\hat{E}(0,t)}{mc\omega_p} \simeq \frac{\bar{\sigma}_J j_0}{2} (\mathbf{i}-1)\sqrt{t}.$$
(5)

Figure 2(a) are snapshots of the diffusing fields at different times, obtained from numerical integration of equation (2). As the field diffuses, its modulation length shrinks with increasing distance for fixed time, while growing at a fixed position as can be predicted from the cosine term in equation (4). This temporal behaviour is presented in figure 2(b), where the result of numerical integration of equation (2) is plotted at $\omega_p x/c = 0$ (current centre), 5, and 10. The central field increases monotonically by \sqrt{t} as expected from equation (5), while



Figure 3. (a) Diffusion of the transverse electromagnetic field inside a magnetized plasma (B = 2 T) and THz radiation with $n_0 = 5 \times 10^{18}$ cm⁻³, $a_0 = 0.05$, and the pulse duration σ ($= c\tau$) = 4 μ m. The snapshot was captured at 1.3 ps after the pulse collision. (b) The field measured on the axis. (c) Temporal growth and saturation of the THz field for different collision points (L), and (d) the corresponding spectra of the THz electric field, with $n_0 = 1.25 \times 10^{18}$ cm⁻³, $\sigma = 8 \,\mu$ m (31.4 fs at full-width-half-maximum (FWHM), red, blue), and $n_0 = 5 \times 10^{18}$ cm⁻³, $\sigma = 4 \,\mu$ m (15.7 fs at FWHM, green). The normalization factor of the electric field, $mc\omega_0/e = 3.7$ TV m⁻¹.

the off-central field initially follows $t^{3/2}$ obeying equation (4), but eventually reduces to \sqrt{t} as the central peaked region of the field expands.

The simultaneous diffusion and growth of the field can be utilized as a method to convert the plasma oscillation into an electromagnetic wave in free space. As the growing field eventually hits the plasma-vacuum boundary through diffusion, radiation will be emitted into free space with temporally increasing amplitude. Indeed, this field growth driven by the diffusion mechanism is the major advantage of our scheme in producing strong THz emission. To confirm this scenario of THz radiation, we have performed one- and two-dimensional particle-in-cell (PIC) simulations. A trapezoidal shape has been chosen for the plasma density profile, with two different densities in the flat region, 1.25×10^{18} and 5×10^{18} cm⁻³ (10 and 20 THz, respectively). To reduce any mismatch of radiation impedance, a density ramp-up over 100 μ m was added to the flat plasma. An additional important effect of the density gradient is that the field growth indicated by equation (4) is sustained for a longer time, eventually leading to stronger THz emission. The two counter-propagating pulses are arranged so that they collide at 25 or 50 μ m from the knee of the density gradient. The wavelength of one of the pulses is 870 nm, which is typical for Ti:sapphire lasers, and the other is detuned so that their beat resonantly drives the plasma oscillation. In the two-dimensional simulation, the pulses focus at the colliding point with 50 μ m spot radius. The normalized vector potential of the pulse is 0.05, for which $I \sim 5 \times 10^{15}$ W cm⁻² and $P \sim 0.2$ TW.

Figure 3(a) is an image of the THz emission obtained from two-dimensional PIC simulations. The diffusion and growth of the field measured on the axis, figure 3(b), takes on a very similar shape to the theoretical model. To observe the long-time behaviour of the signals, several one-dimensional PIC simulations have also been performed with the same parameters, varying the distance from the plasma edge to the pulse collision point as shown in figure 3(c). In this figure, THz emission grows as $t^{3/2}$ initially, but soon evolves into a \sqrt{t} dependence, which is exactly the same feature as in figure 2(b). When the pulse collision occurs further into the plasma, it takes longer for the emission to grow, but eventually it reaches a comparable level (red and blue). Note that, due to the density gradient, a strong, but short duration emission by linear mode conversion [10] emerges simultaneously in the early stage. Though not fully plotted in the figure, the emission usually lasts up to an order of a hundred pico-seconds, which produces quite a monochromatic frequency spectrum as shown in figure 3(d). In the PIC simulations, collisions between charged particles are neglected, since the collisional rate is very small compared with the plasma frequency for the given density. Furthermore, we did not consider collisions between charged and neutral particles, assuming the plasma is prepared one hundred percent ionized by discharge or field ionization. If the ionization is not complete, collisions can reduce the lifetime of the current oscillation, leading to a shortened THz pulse.

3. Scaling of the THz amplitude

To obtain a scaling law for the field strength of the THz emission as a function of the driving pulse amplitude, we calculate the radiation current source, i.e. the J_0 term in equation (2). Since the current is driven by the beat of two counter-propagating pulses, the electron oscillation is spatially fast-varying, so that a linear portion of the oscillation is averaged out. However, the remaining smooth, nonlinear component can still be strong [31], forming a bunched longitudinal current oscillating at the plasma frequency. Once this average longitudinal current is generated, it induces a transverse current oscillation via the external magnetic field.

If the right- and left-going laser pulses are represented by $a_{+} = \hat{a}_{+}e^{ik_{+}z-i\omega_{+}t}$ and $a_{-} = \hat{a}_{-}e^{-ik_{-}z-i\omega_{-}t}$, respectively, their beat is proportional to $e^{i(k_{+}+k_{-})z-i\Delta\omega t}$. Thus, they exert a ponderomotive force on the electrons which yields a density perturbation $n_{1} = \frac{1}{2}(\hat{n}_{1}e^{i(k_{+}+k_{-})z-i\Delta\omega t} + c.c.)$. Because the spatial average of n_{1} disappears, the linear current $-en_{0}v_{1}$ does not contribute significantly to the electromagnetic radiation. However, a non-vanishing second-order current remains, given by

$$\langle J_2 \rangle = -\frac{e}{4} \left(\hat{n}_1 \hat{v}_1^* + \hat{n}_1^* \hat{v}_1 \right) = -\frac{e\omega_p}{8kn_0} \left| \hat{n}_1 \right|^2.$$
 (6)

Here, we have used the continuity relation between n_1 and v_1 , and assumed resonant driving, i.e. $\Delta \omega \simeq \omega_p$. This spatially smooth nonlinear current, together with the linear self-current, drives the longitudinal electric field oscillations via the relation $\partial_t E_{z,\text{smooth}} = -4\pi (J_{1,\text{self}} + \langle J_2 \rangle)$. Calculating the self-current using the linearized equation of motion for the electron plasma, we obtain

$$\left(\frac{\partial^2}{\partial t^2} + \omega_p^2\right) E_{z,\text{smooth}} = \frac{\pi e \omega_p}{2kn_0} \frac{\partial}{\partial t} \left| \hat{n}_1 \right|^2.$$
(7)

Note that $k_+ \simeq k_- = k$, so $k_+ + k_- \simeq 2k$. To obtain the solution of equation (7) we evaluate the temporal evolution of \hat{n}_1 driven by the beat of the counter-propagating pulses

$$\frac{\partial \hat{n}_1}{\partial t} = -i \frac{n_0 c^2 k^2}{\omega_p} \hat{a}_+ \hat{a}_-^*,\tag{8}$$

where n_0 represents the unperturbed plasma density. In deriving equation (8), we neglect the magnetic force term, because we consider a very weakly magnetized plasma, i.e. $\omega_c \ll \omega_p$. In the case where the plasma oscillation is driven by the beat of counter-propagating pulses, as in equation (8), the transverse component of the ponderomotive force is commonly neglected [31]. The transverse force is proportional to a^2/w^2 , where *w* is the pulse spot size, while the longitudinal one by the counter-beat is $\sim k^2 a^2$. Since $k \gg 1/w$ under usual circumstances, the longitudinal ponderomotive force is dominant. For laser pulses assumed to be longitudinally Gaussian with equal maximum amplitude a_0 and pulse duration τ , the pulse amplitudes can be written as $\hat{a}_{\pm} = a_0 \exp\left[-(t \mp (z \pm L)/c)^2/\tau^2\right]$. Note that it is arranged for the peaks of the pulses to overlap at z = 0 after propagation by distance *L*. Then, applying the Green function method to equation (7), we obtain

$$E_{z,\text{smooth}} = A(z) \int_{-\infty}^{\infty} e^{i\omega_p \left[t - \left(\xi \tau / \sqrt{2} + L/c\right) \right]} \Phi(\xi) \, \mathrm{d}\xi, \tag{9}$$

where

$$A(z) = \pi^{2/3} \frac{e n_0 c^4 k^3 a_0^4 \tau^2}{2i\omega_p^2} \exp\left(-4z^2 / \tau^2 c^2\right)$$

and $\Phi(\xi) = \exp(-\xi^2)(1 + \operatorname{erf}(\xi))/2$. Finally, the spatially averaged amplitude of the longitudinal oscillation becomes

$$\frac{eE_{z,\text{smooth}}}{mc\omega_0} = \frac{\pi}{16} \sigma^2 k^2 a_0^4 e^{-0.084\omega_p^2 \tau^2} e^{-4z^2/\sigma^2},$$
(10)

where ω_0 is the laser frequency and $\sigma = c\tau$ is the spatial length of the laser pulse. An interesting feature of this equation is that the longitudinal field depends only weakly on the plasma density via the exponential term, which is quite different from the single-pulse-driven wakefield.

By applying an external magnetic field in the transverse direction, the longitudinal electric field given by equation (10) can be partially converted to a transverse field oscillating with the same frequency ω_p . Here the ratio of the transverse oscillation to the longitudinal one is given by $E_y = \omega_c / \omega_p E_{z,\text{smooth}}$, where $\omega_c = eB_0/m$, leading to





$$\frac{eE_y}{mc\omega_0} = \frac{\pi}{16} \frac{\omega_c}{\omega_p} \sigma^2 k^2 a_0^4 e^{-0.084\omega_p^2 \tau^2}.$$
 (11)

The radiation source current is then calculated from the relation $J_{y} = -\varepsilon_{0}\partial_{t}E_{y}$.

If the plasma-vacuum boundary is sharp, and the pulse collision point is located exactly at the edge of the plasma, the transverse field given by equation (11) is coupled to the vacuum without experiencing the diffusion and growth. Thus equation (11) gives the scaling of the minimum THz amplitude as a function of the driving pulse amplitude. An interesting point in this scaling is that the THz amplitude is proportional to a_0^4 , corresponding to P^2 , where *P* is the power of the driving laser pulse. This is significantly different from the single-pulse driven systems, where the emission amplitude is proportional just to P [30]. From this different scaling, and due to the k^2 -factor, one of which is replaced by a much smaller value σ^{-1} in the single-pulse case, we expect the counter-propagating pulse scheme to generate a much stronger THz wave than the single-pulse driven systems with very moderate laser power. More detailed comparison between equation (11) and the single-pulse scaling shows that the effect of colliding pulses dominates the single-pulse effect when $a_0 > a_{th}$, where

$$a_{\rm th} \sim 0.42 \frac{1}{\sqrt{N}} \frac{\omega_p}{\omega_0}.$$
 (12)

Here *N*, which represents the number of oscillations of the laser field within the pulse duration, is usually >10, and $\omega_p/\omega \ll 1$, thus this condition is satisfied even for small a_0 .

Figure 4 shows theoretical curves of the THz amplitudes from the two different schemes for $n_0 = 5 \times 10^{18}$ and $n_0 = 1.25 \times 10^{18}$ cm⁻³, corresponding to 20 and 10 THz, respectively.

In figure 4, the simulation data (circles) exhibits saturation, which arises from wave-breaking of the spatially fast varying wave. One major effect of wave-breaking is the suppression of the plasma current by kinetic detuning [32]. From the fact that it occurs when the electron fluid velocity, which is $\hat{v}_1 \simeq \sqrt{\pi/8} a_0^2 \omega \sigma/2$ according to the linearized continuity equation, exceeds the phase velocity of the plasma wave, we estimate the laser amplitude for saturation to be

$$a_{\rm sat} = \left(\frac{8}{\pi}\right)^{1/4} \sqrt{\frac{\omega_p}{\omega_0 \sigma k}}.$$
 (13)

In addition, though not as apparent in the figure, the scaling for very low a_0 , less than a_{th} , obeys a_0^2 rather than a_0^4 . As a consequence, operating with a_0 between a_{th} and a_{sat} may yield optimum energy conversion from laser to THz emission.

As mentioned above in the discussion of figure 3(b), the density gradient helps the diffusing field to grow and to yield much stronger THz emission than for the sharp boundary case. This feature is verified by the PIC simulations, as shown in figure 5, where a significant enhancement due to the density gradient is apparent. Indeed, the positive effect of the density gradient is one of the advantages of the field diffusion mechanism. Because every realistic plasma from a gas-jet or capillary discharge used for laser-plasma interactions has a natural density ramp-up, the experimental conditions can be greatly relaxed in our diffusion-growth scheme. Furthermore, even though a varying density is employed, the radiation frequency is not influenced much by that since our scheme is based on the local oscillation of the plasma. Note that for a density gradient the THz frequency is usually chirped when driven by a single pulse, leading to broad-band emission.



Figure 5. Dependence of the amplitude of THz emission on the length of the density gradient for $n_0 = 1.25 \times 10^{18}$ cm⁻³ (10 THz), $a_0 = 0.05$, $\sigma = 8 \mu$ m, and B = 2 T.

4. Conclusions

In summary, we have proposed and investigated a novel method for obtaining strong THz emission based on a driven-diffusion mechanism. We obtain, analytically, the temporal growth of the diffusing electromagnetic field by solving the constantly driven complex diffusion equation. For the driving term of the diffusing field a long lasting localized current source for the radiation can be produced by two counter-propagating laser pulses colliding in a weakly magnetized plasma. From analytic theory, we find that the THz amplitude scales with P^2 , where P is the power of the driving pulse, which is verified by one-dimensional PIC simulations. This scaling gives much stronger emission than for single-pulse driven systems, for which it is proportional just to P. Such significant enhancement of the THz amplitude arises from two factors: one is the stronger ponderomotive force produced by counter-propagating pulses compared with a single pulse for a given total energy of the pulses. The other is the growth of the emission by a driven-diffusion mechanism of the electromagnetic field.

Finally, we have confirmed the feasibility of the proposed scheme for experimental realization by showing that it does *not* impose any strict condition on the plasma or lasers such as a sharp plasma edge or the exact pulse collision point. Moreover, we have discovered that the density gradient positively enhances the emission amplitude. From these results, we note that the experimental conditions can be significantly relaxed. We suggest the following proof-of-principle experimental parameters: a couple of millimetres plasma with density of order 10^{18} cm⁻³ and density-ramp-up over ~ 100μ m, which is readily available from gas jet or capillary discharge. The intensity of the laser pulses with 30 fs FWHM pulse duration used in our simulation is 5×10^{15} W cm⁻², which yields a THz amplitude of 50 MV m⁻¹. Therefore we expect that pulses with a few TW power and 200 μ m spot radius (pulse energy ~0.2 J) may yield a THz power of order 0.1 MW.

Acknowledgments

This research was supported by the Basic Science Research Program through the National Research Foundation (NRF) of Korea funded by the Ministry of Science, ICT and Future Planning (Grant number NRF-2014M1A7A1A01030175 and NRF-2013R1A1A2006353, NRF-2014M1A7A1A01030173). We acknowledge the support of the UK EPSRC (grant no. EP/J018171/1), the ECs LASERLAB-EUROPE (grant agreement no. 284464, Seventh Framework Programme), EuCARD-2 (grant no. 312453, FP7) and the Extreme Light Infrastructure (ELI) European Project. We greatly appreciate Prof Z -M Sheng for useful comments on our work.

References

- Hamster H, Sullivan A, Gordon S, White W and Falcone R W 1993 Subpicosecond, electromagnetic pulses from intense laser-plasma interaction *Phys. Rev. Lett.* 71 2725
- [2] Hamster H, Sullivan A, Gordon S and Falcone R W 1994 Short-pulse terahertz radiation from high-intensity-laser-produced plasmas Phys. Rev. E 49 671
- [3] Cheng C-C, Wright E M and Moloney J V 2001 Generation of electromagnetic pulses from plasma channels induced by femtosecond light strings Phys. Rev. Lett. 87 213001
- [4] Cheng C-C, Wright E M and Moloney J V 2002 Cheng, Wright Moloney reply Phys. Rev. Lett. 89 139302
- [5] Shvets G, Kaganovich I and Startsev E 2002 Comment on generation of electromagnetic pulses from plasma channels induced by femtosecond light strings Phys. Rev. Lett. 89 139301
- [6] Cao L-H, Yu W, Xu H, Zheng C-Y, Liu Z-J, Li B and Bogaerts A 2004 Terahertz radiation from oscillating electrons in laser-induced wake fields Phys. Rev. E 70 046408

- [7] Sprangle P, Penano J R, Hafizi B and Kapetanakos C A 2004 Ultrashort laser pulses and electromagnetic pulse generation in air and on dielectric surfaces Phys. Rev. E 69 066415
- [8] Wu H-C and Meyer-ter-Vehn J 2009 Terahertz radiation from a laser plasma filament 36th EPS Conf. Plasma Phys. Sofia, ECA vol 33E, P-2.027
- [9] Sheng Z-M, Mima K, Zhang J and Sanuki H 2005 Emission of electromagnetic pulses from laser wakefields through linear mode conversion *Phys. Rev. Lett.* 94 095003
- [10] Wu H-C, Sheng Z-M, Dong Q-L, Xu H and Zhang J 2007 Powerful terahertz emission from laser wakefields in inhomogeneous magnetized plasmas Phys. Rev. E 75 016407
- [11] Lőffler T, Jacob F and Roskos H G 2000 Generation of terahertz pulses by photoionization of electrically biased air Appl. Phys. Lett. 77 453
- [12] Cook D J and Hochstrasser R M 2000 Intense terahertz pulses by four-wave rectification in air Opt. Lett. 25 1210
- [13] Xie X, Dai J and Zhang X C 2006 Coherent control of THz wave generation in ambient air Phys. Rev. Lett. 96 075005
- [14] Kim K Y, Glownia J H, Taylor A J and Rodriguez G 2007 Terahertz emission from ultrafast ionizing air in symmetry-broken laser fields Opt. Express 15 4577
- [15] Kim K Y, Taylor A J, Glownia J H and Rodriguez G 2008 Coherent control of terahertz supercontinuum generation in ultrafast lasergas interactions *Nat. Photonics* 2 605
- [16] Clerici M et al 2013 Wavelength scaling of terahertz generation by gas ionization Phys. Rev. Lett. 110 253901
- [17] Vvedenskii N V, Korytin A I, Kostin V A, Murzanev A A, Silaev A A and Stepanov A N 2014 Two-color laser-plasma generation of terahertz radiation using a frequency-tunable half harmonic of a femtosecond pulse *Phys. Rev. Lett.* **112** 055004
- [18] Yoshii J, Lai C H, Katsouleas T, Joshi C and Mori W B 1997 Radiation from Cerenkov wakes in a magnetized plasma Phys. Rev. Lett. 79 4194
- [19] Yugami N, Higashiguchi T, Gao H, Sakai S, Takahashi K, Ito H and Nishida Y 2002 Experimental observation of radiation from Cherenkov wakes in a magnetized plasma *Phys. Rev. Lett.* 89 065003
- [20] Dorranian D, Starodubtsev M, Kawakami H, Ito H, Yugami N and Nishida Y 2003 Radiation from high-intensity ultrashort-laserpulse and gas-jet magnetized plasma interaction *Phys. Rev. E* 68 026409
- [21] Bergé L, Skupin S, Köhler C, Babushkin I and Herrmann J 2013 3D numerical simulations of thz generation by two-color laser filaments Phys. Rev. Lett. 110 073901
- [22] Babushkin I, Kuehn W, Köhler C, Skupin S, Bergé L, Reimann K, Woerner M, Herrmann J and Elsaesser T 2010 Ultrafast spatiotemporal dynamics of terahertz generation by ionizing two-color femtosecond pulses in gases Phys. Rev. Lett. 105 053903
- [23] Babushkin I, Skupin S, Husakou A, Köhler C, Cabrera-Granado E, Bergé L and Herrmann J 2011 Tailoring terahertz radiation by controlling tunnel photoionization events in gases New J. Phys. 13 123029
- [24] Weling A S, Hu B B, Froberg N M and Auston D H 1994 Generation of tunable narrow-band THz radiation from large aperture photoconducting antennas Appl. Phys. Lett. 64 137
- [25] Liu Y, Park SG and Weiner A M 1996 Enhancement of narrow-band terahertz radiation from photoconducting antennas by optical pulse shaping Opt. Lett. 21 1762
- [26] Dobroiu A, Otani C and Kawase K 2006 Terahertz-wave sources and imaging applications Meas. Sci. Technol. 17 R161
- [27] Boboc A, Zabeo L, Murari A and JET EFDA Contributors 2006 Simultaneous Cotton–Mouton and Faraday rotation angle measurements on JET Rev. Sci. Instrum. 77 10F324
- [28] Vieux G, Ersfeld B, Farmer J P, Hur M S, Issac R C and Jaroszynski D A 2013 Plasma density measurements using chirped pulse broadband Raman amplification *Appl. Phys. Lett.* **103** 121106
- [29] Debayle A, Gremillet L, Bergé L and Köhler C 2014 Analytical model for THz emissions induced by laser-gas interaction Opt. Express 22 013691
- [30] Spence N, Katsouleas T, Muggli P, Mori W B and Hemker R 2001 Simulations of Cerenkov wake radiation sources Phys. Plasmas 8 4995
- [31] Shvets G, Fisch N J, Pukhov A and Meyer-ter-Vehn J 1999 Generation of periodic accelerating structures in plasma by colliding laser pulses *Phys. Rev.* E 60 2218
- [32] Hur M S, Lindberg R R, Charman A E, Wurtele J S and Suk H 2005 Electron kinetic effects on Raman backscatter in plasmas Phys. Rev. Lett. 95 115003